


TEXAS CHILDREN'S HOSPITAL EVIDENCE-BASED OUTCOMES CENTER Intraoperative Spinal Instrumentation Prevention Evidence-Informed Pathway



Version 11/2021

Clinical standards are developed for 80% of the patient population with a particular disease. Each practitioner must use his/her clinical judgment in the management of any specific patient.

Critical Points of Evidence***Evidence Supports**

- Administer cefazolin at a dose of 30 mg/kg up to a maximum dose of 2 grams for patients under 120 kg, and 30 mg/kg up to a maximum dose of 3 grams for patients over 120 kg. Re-dose cefazolin every 4 hours times 2 doses, then every 8 hours (1-3) — Strong recommendation, very low quality evidence
- Irrigation of the surgical site with normal saline. (23-26)
- Use 1-2 grams of vancomycin powder sprinkled in the surgical wound prior to closure to prevent surgical site infection in children that weigh >25kg. (4-10) — Strong recommendation, very low quality evidence
- To provide coverage for gram negative pathogens for incontinent pediatric patients by administering gentamicin at a dosing of 2.5 mg/kg with a maximum dose of 120 mg. (2, 11-14) — Strong recommendation, very low quality evidence
- To use drains in pediatric patients undergoing spinal instrumentation surgery for scoliosis. (3, 15-17) — Strong recommendation, very low quality evidence

Evidence Lacking/Inconclusive

- Irrigation by hand compared to pulse irrigation to decrease surgical site infection. (13, 18-20) — Unable to make a recommendation
- Use a three-minute povidone-iodine soak decrease the risk of surgical site infection. (21-22) — Unable to make a recommendation
- To administer vancomycin as the prophylactic pre-incision antibiotic in patients with severe penicillin allergy (defined by anaphylaxis) or known cephalosporin allergy. — Consensus recommendation

*NOTE: The references cited represent the entire body of evidence reviewed to make each recommendation.

Measures**Process**

- Timely prophylactic antibiotic administration
- Maintenance of normothermia
- Appropriate prophylactic antibiotic re-dosing

Outcome

- Mortality
- Morbidity
- Instrumented spine surgical site infections
- Length of stay

References

1. Cies, J. J., Chan, S., Hossain, J., Brenn, B. R., & Di Pentima, M. C. (2012). Influence of body mass index and antibiotic dose on the risk of surgical site infections in pediatric clean orthopedic surgery. *Surgical Infections*, 13(6), 371-376.
2. Linam, W. M., Margolis, P. A., Staat, M. A., Britto, M. T., Hornung, R., Cassidy, A., & Connelly, B. L. (2009). Risk factors associated with surgical site infection after pediatric posterior spinal fusion procedure. *Infection Control & Hospital Epidemiology*, 30(2), 109-116.
3. Ramo, B. A., Roberts, D. W., Tuason, D., McClung, A., Paraisson, L. E., Moore, H. G. 4th, & Sucato, D. J. (2014). Surgical site infections after posterior spinal fusion for neuromuscular scoliosis: A thirty-year experience at a single institution. *Journal of Bone and Joint Surgery America*, 96(24), 2038-2048.
4. Tubaki, V. R., Rajasekaran, S., & Shetty, A. P. (2013). Effects of using intravenous antibiotic only versus local intrawound vancomycin antibiotic powder application in addition to intravenous antibiotics on postoperative infection in spine surgery in 907 patients. *Spine (Phila Pa 1976)*, 38(25), 2149-2155.
5. Gans, I., Dormans, J. P., Spiegel, D. A., Flynn, J. M., Sankar, W. N., Campbell, R. M., & Baldwin, K. D. (2013). Adjunctive vancomycin powder in pediatric spine surgery is safe. *Spine (Phila Pa 1976)*, 38(19), 1703-1707.
6. Ghobrial, G. M., Thakkar, V., Andrews, E., Lang, M., Chitale, A., Oppenlander, M. E., et al. (2014). Intraoperative vancomycin use in spinal surgery: Single institution experience and microbial trends. *Spine (Phila Pa 1976)*, 39(7), 550-555.
7. Molinari, R. W., Khera, O. A., & Molinari, W. J., 3rd. (2012). Prophylactic intraoperative powdered vancomycin and postoperative deep spinal wound infection: 1,512 consecutive surgical cases over a 6-year period. *European Spine Journal*, 21 Suppl 4, S476-482.
8. O'Neill, K. R., Smith, J. G., Abtahi, A. M., Archer, K. R., Spengler, D. M., McGirt, M. J., & Devin, C. J. (2011). Reduced surgical site infections in patients undergoing posterior spinal stabilization of traumatic injuries using vancomycin powder. *Spine Journal*, 11(7), 641-646.
9. Sweet, F. A., Roh, M., & Sliva, C. (2011). Intrawound application of vancomycin for prophylaxis in instrumented thoracolumbar fusions: Efficacy, drug levels, and patient outcomes. *Spine (Phila Pa 1976)*, 36(24), 2084-2088.
10. Theologis, A. A., Demirkiran, G., Callahan, M., Pekmezci, M., Ames, C., & Deviren, V. (2014). Local intrawound vancomycin powder decreases the risk of surgical site infections in complex adult deformity reconstruction: A cost analysis. *Spine (Phila Pa 1976)*, 39(22), 1875-1880.
11. Labbe, A. C., Demers, A. M., Rodrigues, R., Arlet, V., Tanguay, K., & Moore, D. L. (2003). Surgical-site infection following spinal fusion: A case-control study in a children's hospital. *Infection Control & Hospital Epidemiology*, 24(8), 591-595.
12. McLeod, L. M., Keren, R., Gerber, J., French, B., Song, L., Sampson, N. R., et al. (2013). Perioperative antibiotic use for spinal surgery procedures in US children's hospitals. *Spine (Phila Pa 1976)*, 38(7), 609-616.
13. Myung, K. S., Glassman, D. M., Tolo, V. T., & Skaggs, D. L. (2014). Simple steps to minimize spine infections in adolescent idiopathic scoliosis. *Journal of Pediatric Orthopaedics*, 34(1), 29-33.
14. Nunez-Pereira, S., Pellise, F., Rodriguez-Pardo, D., Pigrau, C., Sanchez, J. M., Bago, J., et al. (2011). Individualized antibiotic prophylaxis reduces surgical site infections by gram-negative bacteria in instrumented spinal surgery. *European Spine Journal*, 20 Suppl 3, 397-402.
15. Glennie et al. (2015). Dressings and drains in posterior spine surgery and their effect on wound complications.
16. Ho, C., Sucato, D. J., & Richards, B. S. (2007). Risk factors for the development of delayed infections following posterior spinal fusion and instrumentation in adolescent idiopathic scoliosis patients. *Spine (Phila Pa 1976)*, 32(20), 2272-2277.
17. Olsen, M. A., Nepple, J. J., Riew, K. D., Lenke, L. G., Bridwell, K. H., Mayfield, J., & Fraser, V. J. (2008). Risk factors for surgical site infection following orthopaedic spinal operations. *Journal of Bone and Joint Surgery America*, 90(1), 62-69.
18. Nikfarjam, M., Weinberg, L., Fink, M. A., Muralidharan, V., Starkey, G., Jones, R., et al. (2014). Pressurized pulse irrigation with saline reduces surgical-site infections following major hepatobiliary and pancreatic surgery: Randomized controlled trial. *World Journal of Surgery*, 38(2), 447-455.
19. Hayashi, T., Shirane, R., Yokosawa, M., Kimiwada, T., & Tominaga, T. (2010). Efficacy of intraoperative irrigation with saline for preventing shunt infection. *Journal of Neurosurgery: Pediatrics*, 6(3), 273-276.
20. Watanabe, M., Sakai, D., Matsuyama, D., Yamamoto, Y., Sato, M., & Mochida, J. (2010). Risk factors for surgical site infection following spine surgery: Efficacy of intraoperative saline irrigation. *Journal of Neurosurgery: Spine*, 12(5), 540-546.
21. Chang, F. Y., Chang, M. C., Wang, S. T., Yu, W. K., Liu, C. L., & Chen, T. H. (2006). Can povidone-iodine solution be used safely in a spinal surgery? *European Spine Journal*, 15(6), 1005-1014.
22. Cheng, M. T., Chang, M. C., Wang, S. T., Yu, W. K., Liu, C. L., & Chen, T. H. (2005). Efficacy of dilute betadine solution irrigation in the prevention of postoperative infection of spinal surgery. *Spine (Phila Pa 1976)*, 30(15), 1689-1693.
23. Lemans, J. V., Wijdicks, S. P., Boot, W., Govaert, G. A., Houwert, R. M., Öner, F. C., & Kruyt, M. C. (2019). Intrawound treatment for prevention of surgical site infections in instrumented spinal surgery: a systematic comparative effectiveness review and meta-analysis. *Global spine journal*, 9(2), 219-230.
24. Cohen, L. L., Schwend, R. M., Flynn, J. M., Hedequist, D. J., Karlin, L. I., Emans, J. B., ... & Glotzbecker, M. P. (2020). Why irrigate for the same contamination rate: wound contamination in pediatric spinal surgery using betadine versus saline. *Journal of Pediatric Orthopaedics*, 40(10), e994-e998.
25. do Nascimento, T. L., Finger, G., Sfreddo, E., de Lima Cecchini, A. M., de Lima Cecchini, F. M., & Stefani, M. A. (2019). Double-blind randomized clinical trial of vancomycin in spinal arthrodesis: no effects on surgical site infection. *Journal of Neurosurgery: Spine*, 32(3), 473-480.
26. Cannon, J. G., Ho, A. L., Mohole, J., Pendharkar, A. V., Sussman, E. S., Cheshier, S. H., & Grant, G. A. (2019). Topical vancomycin for surgical prophylaxis in non-instrumented pediatric spinal surgeries. *Child's Nervous System*, 35(1), 107-111.

Clinical Standards Preparation

This clinical standard was prepared by the Evidence-Based Outcomes Center (EBOC) team in collaboration with content experts at Texas Children's Hospital. Development of this clinical standard supports the TCH Quality and Patient Safety Program initiative to promote clinical standards and outcomes that build a culture of quality and safety within the organization.

Intraoperative Spinal Instrumentation Infection Prevention Content Expert Team

Melissa Ard, RN, Clinical Data Specialist
Judith Campbell, MD, Infectious Disease
Joyce Enoch, RN, Director of System Accreditation
Darrell Hanson, MD, Orthopedic Surgery
Kenneth Kocab, RN, Outcomes and Impacts Service
Lucila Marquez, MD, Infectious Disease
Laura Monson, MD, Plastic Surgery
Debra Palazzi, MD, Infectious Disease
Nihar Patel, MD, Anesthesiology
Imelda Tjia, MD, Anesthesiology
Veronica Velez, RN, Surgery
Elaine Whaley, MSN, RN, Infection Control

EBOC Team

Sheesha Porter, MSN, RN, Evidence-Based Practice Specialist
Kristy Rialon, MD, Surgery Quality
Binita Patel, MD, Chief Medical Quality Officer

Additional EBOC Support

Andrea Jackson, MBA, RN, Evidence-Based Practice Specialist
Betsy Lewis, MSN, RN, CNL, Evidence-Based Practice Specialist
Anne Dykes, MSN, RN, Manager

No relevant financial or intellectual conflicts to report.

Development Process

This clinical standard was developed using the process outlined in the EBOC Manual. The literature appraisal documents the following steps:

- Review Preparation
 - PICO questions established
 - Evidence search confirmed with content experts
- Review of Existing External Guidelines
 - Clinical Practice Guidelines for Antimicrobial Prophylaxis in Surgery, 2013
 - National Institute for Clinical Excellence, Surgical Site Infection: Prevention and Treatment of Surgical Site Infection, 2020
 - Children's Hospital of Philadelphia, Spine Deformity "Best Practice" Guidelines, 2013
 - Seattle Children's Spine Pathway, 2019
 - Children's Hospital Colorado, High-Risk Spinal Fusion Clinical Care Guideline, 2018
 - Children's Hospital of Orange County, Spinal Fusion for Adolescent Idiopathic Scoliosis Care Guideline, 2018
 - Cincinnati Children's Hospital, Spine Surgical Site Infection Prevention Protocol, 2014
- Literature Review of Relevant Evidence
 - Searched: PubMed, Cochrane Collaboration, Google
- Critically Analyze the Evidence
 - 2 meta-analyses, 6 randomized **controlled** trials, and 21 nonrandomized studies
- Summarize the Evidence
 - Materials used in the development of the clinical standard, literature appraisal, and any order sets are maintained in an Intraoperative Spinal Instrumentation Infection evidence-informed review manual within EBOC.

Evaluating the Quality of the Evidence

Published clinical guidelines were evaluated for this review using the **AGREE II** criteria. The summary of these guidelines are included in the

literature appraisal. AGREE II criteria evaluate Guideline Scope and Purpose, Stakeholder Involvement, Rigor of Development, Clarity and Presentation, Applicability, and Editorial Independence using a 4-point Likert scale. The higher the score, the more comprehensive the guideline.

This clinical standard specifically summarizes the evidence *in support of* or *against* specific interventions and identifies where evidence is *lacking/inconclusive*. The following categories describe how research findings provide support for treatment interventions.

"Evidence Supports" provides evidence to support an intervention
"Evidence Against" provides evidence against an intervention.

"Evidence Lacking/Inconclusive" indicates there is insufficient evidence to support or refute an intervention and no conclusion can be drawn *from the evidence*.

The **GRADE** criteria were utilized to evaluate the body of evidence used to make practice recommendations. The table below defines how the quality of the evidence is rated and how a strong versus weak recommendation is established. The literature appraisal reflects the critical points of evidence.

Recommendation	
STRONG	Desirable effects clearly outweigh undesirable effects or vice versa
WEAK	Desirable effects closely balanced with undesirable effects
Quality	Type of Evidence
High	Consistent evidence from well-performed RCTs or exceptionally strong evidence from unbiased observational studies
Moderate	Evidence from RCTs with important limitations (e.g., inconsistent results, methodological flaws, indirect evidence, or imprecise results) or unusually strong evidence from unbiased observational studies
Low	Evidence for at least 1 critical outcome from observational studies, RCTs with serious flaws or indirect evidence
Very Low	Evidence for at least 1 critical outcome from unsystematic clinical observations or very indirect evidence

Recommendations

Practice recommendations were directed by the existing evidence and consensus amongst the content experts. Patient and family preferences were included when possible. The Content Expert Team and EBOC team remain aware of the controversies in the diagnosis/management of children undergoing spinal instrumentation surgery for scoliosis. When evidence is lacking, options in care are provided in the clinical standard and the accompanying order sets (if applicable).

Approval Process

Clinical standards are reviewed and approved by hospital committees as deemed appropriate for its intended use. Clinical standards are reviewed as necessary within EBOC at Texas Children's Hospital. Content Expert Teams are involved with every review and update.

Disclaimer

Practice recommendations are based upon the evidence available at the time the clinical standard was developed. Clinical standards (guidelines, summaries, or pathways) **do not** set out the standard of care and are not intended to be used to dictate a course of care. Each physician/practitioner must use his or her independent judgment in the management of any specific patient and is responsible, in consultation with the patient and/or the patient's family, to make the ultimate judgment regarding care.

Version History

Date	Comments
Jun 2016	Originally completed
Jan 2022	recommendation & algorithm update