

OPERATIVE TIMES WITH ROBOTIC ASSISTED LAPAROSCOPIC PYELOPLASTY ARE ASSOCIATED WITH SURGEON EXPERIENCE LEVEL AND TRAINEE LEVEL

Park, Bridget S¹, Soo-Jeong Kim², Hannah A Bachtel², Chester J Koh²

¹ Baylor College of Medicine, Department of Other

² Baylor College of Medicine; Texas Children's Hospital, Urology, Pediatric Urology

Keywords: Robotic surgery, pediatric urology, minimally invasive surgery, Uretero-pelvic junction obstruction

Background: Robotic-assisted laparoscopic pyeloplasty (RALP) is the most commonly performed robotic procedure in pediatric urology. While the safety and efficacy of RALP has been demonstrated there is a paucity of data regarding procedure times, potentially reflecting quality outcomes and costs. We aim to examine surgeon and patient factors associated with pediatric RALP procedure times.

Materials/Methods: Prospectively collected quality improvement data was reviewed for all pediatric patients undergoing RALP at a tertiary care children's hospital (2013 – 2020). Descriptive statistics were utilized to compare total procedure and console times among cohorts.

Results: Among 285 pediatric patients, patient factors such as younger age at the time of the surgery ($p < 0.005$) and lower BMI (< 0.005) were associated with significantly lower total procedure and console times. Surgeon factors including surgeon experience level and trainee level were associated with significant differences in console time and total procedure time, where higher primary surgeon experience were associated with lower total procedure times ($p < 0.005$) and console times ($p < 0.005$). Of note, cases with fellow assistance had higher total procedure times ($p < 0.005$) and console times ($p < 0.005$), likely reflecting the educational activity time differences at most institutions.

Conclusions: Total procedure times and console times of RALP in children are influenced by surgeon and patient factors, with longer times associated with lower primary surgeon experience levels, higher trainee levels, as well as older patient ages and higher BMI. Consideration of such factors may allow for improved pre-operative allocation of resources including operating room scheduling times and cost analyses.

Images / Graph / Table

Variable	Mean Total Procedure Time (min)	95% Confidence Interval (min)	Total Procedure Time P-value
Patient Age			< 0.005
< 1	157.75	146.35 – 169.15	
1 – 4	157.11	145.86 – 168.38	
4 - 10	181.40	170.02 – 192.78	
> 10	206.29	188.65 – 223.92	
Patient Gender			0.329
Male	177.34	169.38 – 185.28	
Female	175.09	165.54 – 184.63	
BMI			< 0.005
< 19.5	168.44	161.62 – 175.26	
19.5 - 25	210.14	184.20 – 236.08	
> 25	214.91	173.05 – 256.77	
Surgeon Experience at Current institution			< 0.005
< 30 cases	227.70	197.15 – 258.24	
30 – 100 cases	194.72	175.04 – 214.31	
> 100 cases	166.20	160.11 – 172.29	
Trainee Level			< 0.005
Fellow	190.93	180.05 – 201.82	
Resident	168.42	160.82 – 176.02	